



Research Article

Morphology of Sella Turcica in Skeletal Class II Subjects

Suja Ani, Jayan James and Prasanth S.P

Dept: of Orthodontics, Govt: Dental College, Thiruvananthapuram, India

Received date: 6 March 2014; Accepted date: 2 April 2014; Published date: 30 March 2015

Academic Editor: Ramón Fuentes

Correspondence should be addressed to: Suja Ani; sujaanig@yahoo.co.in

Copyright © 2015 Suja Ani, Jayan James and Prasanth S.P. Distributed under Creative Commons CC-BY 4.0

Abstract

A study on the morphology of sella turcica in skeletal Class II subjects in an orthodontic patient population showed 50% of the Class II subjects to have normal sella turcica morphology. The proportion of normal sella turcica was statistically significant among Class I as compared to Class II. 15% of the Class II subjects had bridging of sella turcica morphology and 11% of the Class II showed notching of posterior wall.

Keywords: Sella turcica, morphology of sellae, Class II malocclusion, skeletal Class II.

Introduction

The sella turcica is a saddle-shape depression in the sphenoid bone of the human skull. The anterior border of the sella turcica is represented by the tuberculum sellae and the posterior border by the dorsum sellae. The pituitary gland is surrounded by the sella turcica. Two anterior and two posterior clinoid process project over the pituitary fossa. The anterior clinoid processes are formed by the medial and anterior prolongations of the lesser wing of sphenoid bone, and the posterior clinoid process by the endings of dorsum sellae. During embryological development, the sella turcica is a key point for the migration of the neural crest cells to the fronto nasal and maxillary development fields.

The morphology of sella turcica is of importance in determining the cephalometric position of reference point sella. Variations in sella morphology have been discovered in skeletal Class III subjects¹, cleft subjects², in subjects with severe craniofacial deviations³, in dental anomalies⁴, and in syndromes^{5,6,7,8,9,10,11}. Morphology of sella turcica in skeletal Class III malocclusion has been studied by various authors^{12,13,14,15}. But the morphology of sella turcica in skeletal Class II subjects has not been reported. So this study was undertaken.

Materials and Methods

This was designed as a cross sectional study. The study population included cephalometric records of 100 skeletal Class II subjects and 100 Class I subjects, who

reported to the Dept. Of Orthodontics, Govt. Dental College, Thiruvananthapuram for orthodontic treatment. This study was conducted in November 2013. The inclusion criteria were adults with no history of orthodontic or orthodontic-surgical treatment, no history of cleft lip repair, craniofacial deviations or other syndromes; For Class II subjects the inclusion criteria included ANB angle that should be more than 5 degree. The exclusion criterion was the presence of proximal caries.

A template as per description of Axelsson¹⁶ (2004) was prepared, this included the normal and the 5 variations. The cephalograms were available with patients reporting to the dept. Both Class II and Class I study population included males and females.

Result

42% of Class I and 52% Class II subjects included in the study were males and 58% of Class I and 48% of Class II subjects were females [Table I] (Fig 1). The mean age of study population was 18.69 ± 2.7 years (18.26 for males and 19 years for females) for Class II and 21.12 ± 3.46 years (21.26 for males and 20.75 for females) for Class I subjects [Table 2 and Table 3].

50% of Class II subjects included in the study presented with a normal sella turcica

whereas 71% of the subjects in Class I malocclusion had a normal sella turcica morphology ($p < .05$). 9% of Class II subjects and 11% of Class I subjects had an oblique anterior wall. 4% of Class II subjects and 8% of Class I subjects had a double contour of floor. 11% of Class II subjects and 4% of Class I subjects had a pyramidal shape of dorsum sellae ($p > .05$). 15% of Class II subjects presented with sella turcica bridging whereas only 4% of Class I subjects showed sella turcica bridging; this was statistically significant ($p = .014$). 11% of Class II subjects presented with notching of posterior wall whereas only 2% of Class I subjects included in the study had notching of posterior wall, this is statistically significant ($p < .05$) [Table 4 and Table 5] (Fig 2).

The proportion of oblique anterior wall, double contour of floor and pyramidal shape did not differ between skeletal Class I and skeletal Class II subjects [Table 3].

In gender wise classification, there was not a statistically significant difference in the normal sella turcica morphology between males and females [Table 6].

The data on ANB angle of the study population is presented in Table 7 and Table 8.

Socio Demographic Profile:

Table 1: Gender Wise Distribution

Gender	Class I		Class II	
	Count	Percent	Count	Percent
Male	42	42	52	52
Female	58	58	48	48
Total	100		100	

Table 2: Descriptive Statistics for Age

Group	N	Minimum	Maximum	Mean	Std. Deviation
Class I	100	16	33	21.12	3.462
Class II	100	15	29	18.69	2.718

Table 3: Descriptive Statistics for Age Based on Gender in Each Group

Group	Sex	N	Minimum	Maximum	Mean	Std. Deviation
Class I	Male	42	16	30	21.46	3.269
	Female	58	16	33	20.75	3.658
Class II	Male	52	15	25	18.26	2.153
	Female	48	15	29	19.00	3.044

Table 4: Sella Turcica Morphology Based on Group

ST Morphology	Class I		Class II	
	Count	Percent	Count	Percent
Normal Sella Turcica	71	71	50	50.0
Oblique anterior wall	11	11	9	9
Double contour of floor	8	8	4	4
Sella turcica bridge	4	4	15	15
Notching of posterior wall	2	2	11	11
Pyramidal shape	4	4	11	11

Table 5: Comparison of ST Morphology Based on Group

Sella Turcica Morphology		Class I		Class II		p#
		Count	Percent	Count	Percent	
Normal Sella Turcica	No	29	29	50	50	0.004
	Yes	71	71	50	50	
Oblique anterior wall	No	89	89	91	91	0.814
	Yes	11	11	9	9	
Double contour of floor	No	92	92	96	96	0.373
	Yes	8	8	4	4	
Sella turcica bridge	No	96	96	85	85	0.014
	Yes	4	4	15	15	
Notching of posterior wall	No	98	98	89	89	0.018
	Yes	2	2	11	11	
Pyramidal shape	No	96	96	89	89	0.105
	Yes	4	4	11	11	

#: Fisher's Exact Test

Table 6: Distribution According to Gender



ST Morphology	Class I				Class II			
	Count	M	F	p	Count	M	F	p
Normal Sella Turcica	71	36	35	0.826	50	18	32	0.311
Oblique anterior wall	11	5	6	0.754	9	8	1	0.004
Double contour of floor	8	4	4	1.00	4	1	3	0.637
Sella turcica bridge	4	3	1	0.679	15	7	8	0.719
Notching of posterior wall	2	1	1	1.00	11	2	9	0.113
Pyramidal shape	4	3	1	0.619	11	6	5	0.519

Table 7: ANB Values in Class I Subjects

Class	N	Minimum	Maximum	Mean	Std. Deviation
I	100	1	2	1.55	0.500

Table 8: ANB Values in Class II Subjects

Class II	N	Minimum	Maximum	Mean	Std. Deviation	P Value
males	52	6	12	6.94	1.481	> 0.05
females	48	6	12	7.22	1.501	

Figure 1: Distribution According to Gender**Figure 2: Comparison of ST Morphology Based on Group**

Discussion

This cross sectional study describes the morphology of sella turcica in skeletal Class II and skeletal Class I subjects. The study

sample included adult patients with skeletal Class II malocclusion.

Morphology: Classifications of sella turcica morphology were attempted in literature. Appraisal of sella turcica morphology is a valuable aid in assessing the pathology in

the pituitary gland¹⁷. Abnormal sella turcica morphology was reported in cases with severe craniofacial deviations³, various genetic disorders², syndromes⁵⁻¹¹ and in dental anomalies^{4, 18}. Variations in sella turcica morphology have been reported by different authors. 75% of subjects in Silverman³¹'s study had a normal morphology for sella turcica and remaining 25% had an abnormal morphology. In the present study 50% of the adults with skeletal Class II showed a normal morphology of sella turcica.

Gorden¹⁹ (1922) classified the sella turcica morphology into 3 shapes, the circular, the oval or the flat/saucer shaped, their study sample included children of age ranging from 1 year to 12 years and most of them had either circular or oval shaped sella turcica. Davidoff and Epstein²⁰ (1950) put forward the term J shaped sella while Fournier and Denizet²¹ (1965) used the term omega sella. Teal²² (1977) classified the sella anatomy into round, oval or flat. Bruneton et al²³ (1979) studied the normal variants of sella turcica of adults.

Axelsson et al¹⁶ (2004) classified the sella turcica shapes into six types; normal sella turcica, oblique anterior wall, double contoured sella, sella turcica bridge, irregularity (notching) in the posterior part of the sella and pyramidal shape of the dorsum sellae. They analysed a sample from Oslo University Craniofacial Growth Archive (Norwegian 6-12 years) and reported that 71% of males and 65 % of females presented with a normal sella turcica morphology.

67% of the Saudi subjects in Alkofide¹²'s study; 80% of adults in skeletal Class III malocclusion and 70% of adults in skeletal Class I malocclusion in Iraqi adults in Hadeel et al²⁰'s study; 65% of skeletal Class I patients and 72% of skeletal Class III patients from Islamabad in the Shah et al²¹'s study; and 48% of skeletal Class III and 75 % of skeletal Class I subjects included in Sathyanarayana et al²⁴'s study in South Indian population had a normal sella turcica morphology. In the present study, 50% of skeletal Class II subjects and 71% of skeletal Class I subjects had a normal sella turcica morphology. The proportion of normal sella

turcica morphology was statistically significant among skeletal Class I as compared to skeletal Class II.

18.6% of subjects in Becktor et al³'s study and 16.7% of subjects in Jones et al²⁵'s study showed sella turcica bridging which was more than double the incidence of bridging previously reported.

Busch²⁶, Muller²⁷ and Platzter²⁸ reported the prevalence of sella turcica bridge with a frequency of 1.75 to 6 % in the 'normal' population. Becktor et al³'s and Jones et al²⁵'s study was on patients with severe craniofacial deviations who required combined surgical orthodontic treatment. In the present study 4% of skeletal Class I subjects had sella turcica bridging.

Abdel Kader et al²⁹ (2007) showed significantly higher incidence of sella turcica bridging in subjects from Saudi Arabia with skeletal Class III malocclusion when compared to skeletal Class II and skeletal Class I malocclusion. The present study showed 15% of the Class I subjects with sella turcica bridging.

Marcotty et al¹, in their investigations on Caucasian individuals, found 16.8% of skeletal Class III patients and 9.4% of skeletal Class I patients to have sella turcica bridging. Shah et al¹⁴ reported no incidence of sella turcica bridging in Islamabad orthodontic patients.

In a recent study by Sathyanarayana et al²⁴ in South Indian population, 15% of subjects in skeletal Class III subjects and 5% of skeletal Class I subjects were reported to have sella turcica bridging. In the present study 15% of subjects with skeletal Class II malocclusion had sella turcica bridging (Table 5).

Also many authors had concluded that, the prevalence of sella turcica bridging is higher in subjects with dental anomalies, cleft lip and palate and various other anomalies.

Axelsson et al¹⁶ in their study on subjects from Oslo University Craniofacial Growth Archive reported 23 % of males and 3% of females to have an oblique anterior wall morphology. Sathyanarayana et al²⁴ reported 7% of skeletal Class III subjects

and 3% of skeletal Class I subjects (in South Indian population) and Hadeel et al³⁰ reported 3.3% of skeletal Class III subjects and 4% of skeletal Class I subjects included (Iraqi adult population) to have oblique anterior wall morphology of sella turcica. In the present study 9% of skeletal Class II and 11% of skeletal Class I subjects presented with an oblique anterior wall. The proportion of oblique anterior wall morphology of sella turcica did not differ significantly between skeletal Class I and skeletal Class II malocclusion groups included in the study.

In the study on double contour of floor morphology of sella turcica, Alkofide¹²(study on orthodontic patient population from Saudi Arabia) reported 9.4% of the study sample, Shah et al¹⁴ reported 1.6% of skeletal Class III subjects and 5% of skeletal Class I subjects of study sample (Islamabad), Sathyanarayana et al²⁴ reported 7% of skeletal Class III subjects and 3% of skeletal Class I subjects (South Indian population), Hadeel et al³⁰ reported 10% of skeletal Class III subjects and 14% of skeletal Class I subjects(Iraqi adults), Alkofide¹² reported 8.9% of orthodontic patient population (Saudi Arabia) and Axelsson et al¹⁶ reported 3% of females included in the study (Oslo University Craniofacial Growth Archive) to have double contour of floor. In the present study, 8% of Class I and 4% of Class II presented with double contour of floor.

2.8% of orthodontic population (Saudi Arabia) in Alkofide¹²'s study, 5% of females in Axelsson et al¹⁶'s study, 10% of skeletal Class III subjects and 8.3% of skeletal Class I subjects included in Shah et al¹⁴'s study, 7% of skeletal Class III subjects and 2% of skeletal Class I subjects (South Indian population) in Sathyanarayana et al²⁴'s study, 3.3% of skeletal Class III and 4% of skeletal Class I subjects in Hadeel et al³⁰'s study presented with a pyramidal shape morphology of dorsum sella. In the present study 11% of skeletal Class II and 4% of skeletal Class I subjects presented with a pyramidal shape of sella turcica.

In the present study 11% of skeletal Class II subjects showed notching of the posterior part of sella turcica. 17% of Skeletal Class III

subjects and 12% of Skeletal Class I subjects in Sathyanarayana et al²⁴'s study (South Indian population), 3.3% of skeletal Class III subjects and 6% of skeletal Class I subjects Hadeel et al³⁰'s study, 10% of skeletal Class III subjects and 13.3% of skeletal Class I subjects of Shah et al's study (Islamabad), 11.1% of the orthodontic patient population (Saudi Arabia) in Alkofide¹²'s study and 11% of females (Oslo University Craniofacial Growth Archive) in Axelsson et al¹⁶'s study had irregularities of posterior part of dorsum sella.

Conclusion

The present study showed 71% of Class I and 50% of Class II subjects to have normal sella turcica morphology. The variation in proportion of normal sella turcica was statistically significant among Class I as compared to Class II. Bridging of sella turcica and the notching of posterior wall were seen to be significantly more among Class II subjects than among Class I subjects.

References

1. Meyer-Marcotty, P., Reuther, T. & Stellzig-Eisenhauer, A. "Bridging of the Sella Turcica in Skeletal Class III Subjects," *The European Journal of Orthodontics* (2010); 32(2):148-53.
2. Alkofide, EA. "Sella Turcica Morphology and Dimensions in Cleft Subjects," *Cleft Palate Craniofac Journal* (2008); 45(6):647-53.
3. Becktor, J. P, Einersen, S. & Kjaer, I. "A Sella Turcica Bridge in Subjects with Severe Craniofacial Deviations," *The European Journal of Orthodontics*, (2000); 22(1):69-74.
4. Leonardi, R., Barbato, E., Vichi, M. & Caltabiano, M. "A Sella Turcica Bridge in Subjects with Dental Anomalies," *The European Journal of Orthodontics* (2006); 28(6):580-5.
5. Russell, B. G. & Kjaer, I. "Postnatal Structure of the Sella Turcica in Down Syndrome,". *American Journal of Medical Genetics* (1999); 87(2):183-8.

6. Weinstein, M., Tyrrell, B. & Newton, T. H. "The Sella Turcica in Nelson's Syndrome," *Radiology* (1976); 118(2):363-5.
7. Axelsson, S., Storhaug, K. & Kjaer, I. "Post-Natal Size and Morphology of the Sella Turcica in Williams Syndrome," *The European Journal of Orthodontics* (2004); 26(6):613-21.
8. Childers, N. K. & Wright, J. T. "Dental and Craniofacial Anomalies of Axenfeld-Rieger Syndrome," *Journal of Oral Pathology & Medicine* (1986); 15(10):534-9.
9. Koshino, T., Konno, T. & Ohzeki T. "Bone and Joint Manifestations of Rieger's Syndrome: A Report of a Family," *Journal of Pediatric Orthopaedics* (1989); 9(2):224-30.
10. Meyer-Marcotty, P., Weisschuh, N., Dressler, P., Hartmann, J. & Stellzig-Eisenhauer, A. "Morphology of the Sella Turcica in Axenfeld-Rieger Syndrome With PITX2 Mutation," *Journal of Oral Pathology & Medicine* (2008); 37(8):504-10.
11. Venable, H. P. "Empty Sella Syndrome," *Journal of the National Medical Association* (1977); 69(4):243-7.
12. Alkofide, E. A. "The Shape and Size of the Sella Turcica in Skeletal Class I, Class II, and Class III Saudi Subjects," *The European Journal of Orthodontics* (2007); 29(5):457-63.
13. Cutovic, T., Pavlovic, J. & Kozomara, R. "Analysis of Dimensions of Sella Turcica in Patients with Mandibular Prognathism," *Vojnosanit Pregl* (2008); 65(6):456-61.
14. Mahmood Shah, A., Bashir, U., Tasleemiyas. "The Shape and Size of the Sella Turcica in Skeletal Class I, II & III in Patients Presenting at Islamic International Dental Hospital, Islamabad," *Pakistan Oral & Dental Journal June* (2011); 31(1):104-10.
15. Marsan, G. & Oztas, E. "Incidence of Bridging and Dimensions of Sella Turcica in Class I and III Turkish Adult Female Patients," *World Journal of Orthodontics* (2009); 10(2):99-103.
16. Axelsson, S., Storhaug, K. & Kjaer, I. "Post-Natal Size and Morphology of the Sella Turcica. Longitudinal Cephalometric Standards for Norwegians between 6 and 21 Years of Age," *The European Journal of Orthodontics* (2004); 26(6):597-604.
17. Kjaer, I., Keeling, J. W., Reintoft, I., Nolting, D. & Fischer Hansen, B. "Pituitary Gland and Sella Turcica in Human Trisomy 21 Fetuses Related to Axial Skeletal Development," *American Journal of Medical Genetics* (1998); 80(5):494-500.
18. Leonardi, R., Farella, M. & Cobourne, M. T. "An Association between Sella Turcica Bridging and Dental Transposition," *The European Journal of Orthodontics* (2011); 33(4):461-5.
19. Gordon, M. B. "A Roentgenographic Study of the Sella Turcica in Normal Children," *Endocrinology*, (2013).
20. Epstein, B. S. & Davidoff, L.M. "Advanced Atrophy and Enlargement of the Sella Turcica with Destruction of the Sphenoid," *The American Journal of Roentgenology and Radium Therapy* (1951); 66(6):884-93.
21. Fournier A.M, Denizet D. "Omega-Shaped Sella Turcica," *Marseille Médical* (1965); 102(6):503-9.
22. Teal, J. S. "Radiology of the Adult Sella Turcica," *Bulletin of the Los Angeles Neurological Societies* (1977); 42(3-4):111-74.
23. Bruneton, J. N., Drouillard, J.P., Sabatier, J.C., Elie, G.P. & Tavernier, J.F. "Normal Variants of the Sella Turcica: Comparison of Plain Radiographs and Tomograms in 200 Cases 1," *Radiology* (1979); 131(1):99-104.
24. Sathyanarayana, H.P. Kailasam, V. & Chitharanjan, A. B. "The Size And

- Morphology of Sella Turcica in Different Skeletal Patterns among South Indian Population: A Lateral Cephalometric Study," *Journal of Indian Orthodontic Society* (2013);47(4):266-71.74.
25. Jones, R. M., Faqir, A., Millett, D. T., Moos, K. F. & McHugh, S. "Bridging and Dimensions of Sella Turcica in Subjects Treated by Surgical-Orthodontic Means Or Orthodontics Only," *Angle Orthodontist* (2005); 75(5):714-
26. Busch, W. "Die Morphologie der Sella turcica und ihre Beziehungen zur Hypophyse," *Virchows Archiv* (1951); 320:437-58.
27. Müller, F. "Die Bedeutung der Sellabrücke für das Auge," *Klinische Monatsblätter für Augenheilkunde* (1952); 120: 298-302.
28. Platzer, W. "Zur Anatomie der Sellabrücke und Ihrer Beziehung zur A. Carotis interna," *Fortschritte auf Demgebiet Der Roentgenstrahlen und der Nuklearmedizin* (1957) 87:613-16.
29. Abdel-Kader, HM. "Sella Turcica Bridges in Orthodontic and Orthognathic Surgery Patients. A Retrospective Cephalometric Study," *Australian Orthodontic Journal* (2007); 23(1):30-5.
30. Yassir, Y.A, Yousif , H. A. & Nahidh, M. "Size and Morphology of Sella Turcica in Iraqi Adults, ". *Mustansiria Dental Journal* 30-23:(17);(2010).
31. Silverman, F. N. "Roentgen Standards Fo-Size of the Pituitary Fossa from Infancy through Adolescence," *The American Journal of Roentgenology* (1957); 78(3):451-60.