



Case Report

Endovascular Management of a Periinterventional Crural Pseudoaneurysm – A Case Report and Review of Literature

Florian Lüders, Alfred Doku, Nasser Malyar and Holger Reinecke

Department of Cardiology and Angiology, University Hospital of Muenster, Muenster, Germany

Correspondence should be addressed to: Florian Lüders; Florian.Lueders@ukmuenster.de

Received Date: 4 April 2014; Accepted Date: 12 May 2014; Published Date: 31 May 2014

Academic Editor: Kenei Shimada

Copyright © 2014 Florian Lüders, Alfred Doku, Nasser Malyar and Holger Reinecke. Distributed under Creative Commons CC-BY 3.0

Abstract

Iatrogenic pseudoaneurysms of the anterior tibial artery are infrequent; the optimal treatment strategy is still a subject of debate. We report a case of an anterior tibial artery pseudoaneurysm complicating an endovascular intervention which was managed by transluminal platinum coil embolization and review the current literature. Three platinum coils were placed in the anterior tibial artery endovascularly resulting in the successful embolization and occlusion of the pseudoaneurysm. Endovascular therapy offers an effective and a safe strategy in managing infrapopliteal pseudoaneurysms.

Keywords: Peripheral artery disease, pseudoaneurysm, percutaneous transluminal angioplasty

Introduction

Rising awareness and recognition of the burden of obstructive peripheral arterial disease (PAD) in the health care system has also seen a parallel marked increase in utilization of lower extremity endovascular revascularization procedures [1]. Revascularization of infrapopliteal lesions by means of endovascular approach is evolving to be the standard of care, as it provides in general an excellent clinical outcome with low morbidity and at low procedural costs [2]. However, experiences with the management of the complications following PTA, such as iatrogenic infrapopliteal pseudoaneurysm, are sparse and varied.

Infrapopliteal aneurysms, compared to femoral aneurysms, are rare, and consist mainly of pseudoaneurysms. The pathogenesis of a pseudoaneurysm is characterized by a disruption of the arterial wall with extravasation of blood into the surrounding tissue. This leads to the development of a fibrous sac, which gradually increases in size due to the arterial pressure.

Infrapopliteal pseudoaneurysms could develop from penetrating or blunt injury [3,4], bimalleolar fracture or fracture of the ankle [5], infectious processes of the arteries [6] or connective tissue disorders such as Behcet's disease [7] or as a complication of endovascular procedures, like Fogarty-balloon

thromboembolectomy[8], and rarely as a complication of PTA [9,10].

It is expected that with the increase in the number of infrapopliteal endovascular revascularization procedures, the number of procedural related complications will also increase. Such complications frequently necessitate an endovascular therapeutic management [8] since a surgical approach in such patients with comorbidities is associated with an increased morbidity and delayed wound healing.

Here we report a successful endovascular management of an infrapopliteal pseudoaneurysm in a patient with severe claudication (Rutherford category 3) who underwent PTA because of high-degree stenosis of the anterior tibial artery, and review the existing literature.

Case Report

The patient is a 72-year-old woman with a history of dyslipoproteinemia, arterial hypertension and type II diabetes mellitus. She complained of claudication in the right foot and calf after walking a distance of 20 meters. Duplex ultrasonography showed moderate stenosis of the right popliteal

artery, total occlusion of the proximal right ATA and right tibiofibular tract.

Because of clinically distinct pulmonary arterial hypertension associated with CREST syndrome, coronary 3-vessel artery disease, impaired right ventricular systolic function with severe tricuspid regurgitation and her refusal of a surgical treatment, an endovascular approach for revascularization was chosen. Because of the planned PTA, 300 mg loading dose of Clopidogrel was given in addition to her regular platelet inhibition with Aspirin (acetylsalicylic acid). The pre-procedural angiographical image confirmed the findings of the previous Duplex ultrasonography of distal popliteal artery and lower leg arteries (Figure 1). After successful PTA and stenting of the distal popliteal artery and the right tibiofibular tract, PTA of ATA was performed with a 2.5 mm diameter Pacific balloon with an inflation pressure of 8 bar (nominal pressure) over 2 minutes. Immediately after the PTA an extravasation of contrast agent into the surrounding tissue of mid ATA was visible (Figure 2). Six prolonged PTAs with a Pacific balloon (average 7.5 minutes each) with adjuvant administration of 1000 IE protamine stopped the extravasation of contrast medium. The next day, an ultrasound examination revealed a 36 x 23.




Figure 1: Pre-procedural Angiographical Image of Distal Popliteal Artery and Lower Leg Arteries




Figure 2: Post-interventional Extravasation of Contrast Agent into the Surrounding Tissue of Mid ATA, as an Expression of a Pseudoaneurysm

x 21 mm pulsatile pseudoaneurysm with oscillating flow in the ATA (Figure 3). Because of the high operative risk, an

endovascular approach using coil embolization was preferred.




Figure 3: Ultrasound Examination Revealed a 36 X 23 X 21 Mm Pulsatile Pseudoaneurysm with Oscillating Flow in the ATA

Using the right common femoral artery access in antegrade puncture technique, a 110 mm-long, 4-Fr sheath was percutaneously advanced into the vessel. Angiography showed a good short-term result of the interventions at the popliteal artery and of tibiofibular tract. Digital subtraction angiography confirmed a 5.1 × 2.4 cm three-chambered, pulsatile pseudoaneurysm at the middle third of the right ATA. There was no flow in the ATA distal to the pseudoaneurysm. Branches of the fibular artery showed a good blood supply of the foot.

Three microcoils (Boston interlocking coils) were inserted into the ATA just

proximal to the PSA via a micro-catheter (Figure 4). The immediate angiographic control (Figure 4) and duplexsonographic control the next day confirmed a complete occlusion of the pseudoaneurysm with occlusive signal in the ATA. In view of the successful PTA and stenting of the distal popliteal artery and the right tibiofibular tract, the postcoiling ABI value measured in the right ATP (0.75) was slightly improved compared to the pre-interventional value (0.57); the postcoiling ABI value in the right dorsalispedis artery (0.43) was similar to the pre-interventional value (0.44).




Figure 4: Three Microcoils were Inserted in the ATA Just Proximal to the PSA via a Micro-Catheter. The Immediate Angiographic Control Confirmed a Complete Occlusion of the Pseudoaneurysm

The patient improved clinically, and was discharged from hospital 2 days after the coil embolization.

Discussion

Increasing prevalence of atherosclerotic risk factors, particularly diabetes, with an ageing population has resulted in a dramatic increase prevalence of PAD involving more distal, small-sized arteries. These patients frequently have severe comorbidities with a high perioperative risk precluding them from a surgical approach. On the basis of this, the number of endovascular revascularization of the distal parts of lower limb vasculature is increasing worldwide [2]. Despite accumulating experiences with endovascular interventions in the distal arterial bed, increasing the number of such interventions is expected to result also in increasing the number of periprocedural complications such as formation of an intervention-related PSA [10].

The clinical feature of such PSAs is variable. The spectrum ranges from clinically inapparent pseudoaneurysms to the complete scenario of a compartment syndrome necessitating immediate surgical intervention.

There are no evidence-based recommendations or guidelines as to how such complications should be treated. The management of such PSAs depends also on the clinical scenario. In cases of a small (<2 cm) asymptomatic PSA without a tendency

to enlargement, a conservative approach is justified, as such PSA can occlude spontaneously by thrombosis within the lumen of PSA [11]. While PSA of the common femoral or popliteal arteries can often be treated successfully with ultrasound-guided compression, PSA of the infrapopliteal arteries, particularly arising from the ATA is difficult or almost impossible to compress. The surgical management involves the cut off of blood flow to the PSA, either by proximal and distal ligation, or by primary closure of the inflow or by rerouting the blood flow around this damaged segment of the artery. The bypass could be achieved either by using an autologous vein of the patient or by using a synthetic graft [12]. It must be emphasized that surgical approach in small-sized infrapopliteal arteries are associated with a high rate of re-occlusion and abnormal wound healing.

Ultrasound-guided thrombin injection [13] has also been described as an alternative therapeutic option for PSA. Endovascular options include the insertion of a plain or covered stent [14], and the occlusion of the PSA by means of repeated balloon inflation proximal to the PSA neck [15]. Endovascular coil embolization as described in this case might be another option, when the aforementioned are either not possible or not successful. In our patient, repeated inflation of the Pacific balloon did not lead to an occlusion of the PSA.

Looking in the current literature, we found only few cases of endovascular interventions of infrapopliteal PSA [16, 17, 18]. In our case endovascular coil embolization was appropriate because the peroneal artery was the dominant arterial supply to the foot, and therefore the right ATA could be occluded proximal to the PSA neck. The postprocedural obtained peripheral pressures and the ameliorated clinical status of the patient confirmed that the occlusion of the ATA did not result in an aggravation of the right foot ischemia.

Abbreviations

ATA - anterior tibial artery

PAD - peripheral artery disease

PSA - pseudoaneurysm

PTA - percutaneous transluminal angioplasty

Competing Interests

The author(s) declare that they have no competing interests.

References

- Hirsch, A. T., Haskal, Z. J., Hertzler, N. R., Bakal, C. W., Creager, M. A., Halperin, J. L. et al. (2006). "ACC/AHA 2005 Practice Guidelines for the Management of Patients with Peripheral Arterial Disease (Lower Extremity, Renal, Mesenteric, and Abdominal Aortic): A Collaborative Report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients with Peripheral Arterial Disease): Endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; Transatlantic Inter-Society Consensus; and Vascular Disease Foundation," *Circulation* 113: E463-654.
- Conrad, M. F., Kang, J., Cambria, R. P., Brewster, D. C., Watkins, M. T., Kwolek, C. J. & Lamuraglia, G. M. (2009). "Infrapopliteal Balloon Angioplasty for the Treatment of Chronic Occlusive Disease," *Journal of Vascular Surgery*; 50:799-805.
- Edwards, H., Martin, E. & Nowygrod, R. (1982). "Nonoperative Management of a Traumatic Peroneal Artery False Aneurysm," *Journal of Trauma-Injury Infection & Critical Care*, 22:323-6.
- Skudder, P. A., Gelfand, M. L., Blumenberg, R. M. & Fulco, J. (1999). "Tibial Artery False Aneurysm: Uncommon Result of Blunt Injury Occurring During Athletics," *Annals of Vascular Surgery*, 13:589-91.
- Kurian, J., Pillai, S. C. B., Chapple, D. & Frost, R. A. (2003). "Pseudoaneurysm of Peroneal Artery Following Ankle Fracture," *Foot and Ankle Surgery*, 9:233-5.
- McKee, M. A. & Ballard, J. L. (1999). "Mycotic Aneurysms of the Tibioperoneal Arteries," *Annals of Vascular Surgery*, 13(2):188-90.
- Toyota, N., Kimura, F., Yoshida, S., Mitsui, N., Mochizuki, T., Naito, A. & Ito, K. (1999). "Peroneal Artery Aneurysm Treated by Transcatheter Coil Embolization and Temporary Balloon Occlusion in Behcet's Disease," *CardioVascular and Interventional Radiology*, 22(3):257-9.
- Sadat, U., See, T., Cousins, C., Hayes, P. & Gaunt, M. (2007). "Peroneal Artery Pseudoaneurysm-A Case Report and Literature Review," *BMC Surgery*. Mar 29;7:4. Review.
- Derom, A. (2003). "Ruptured Pseudo-Aneurysm of the Popliteal Artery after Percutaneous Transluminal

- Angioplasty," *Acta Chirurgica Belgica*. Feb; 103(1):102-4.
10. Penugonda, N., Duncan, K. & Schreiber, T. (2008). "Popliteal Artery Pseudoaneurysm Following Foxhollow Atherectomy: A Rare Complication," *The Journal of Invasive Cardiology*. Sep; 20(9):477-8.
 11. Kocakoc, E., Bozgeyik, Z. & Ozgocmen, S. (2003). "Spontaneous Occlusion of a Traumatic False Aneurysm of the Peroneal Artery," *Journal of Ultrasound in Medicine*, 22(12):1391-3.
 12. Ramdass, M. J. (2009). "Traumatic False Aneurysm of the Distal Peroneal Artery with Associated anatomic Anomaly: A Case Report," *Cases Journal*; 2: 8605.
 13. Grewe, P. H., Mugge, A., Germing, A. & Harrer, E. (2004). "Occlusion of Pseudoaneurysms Using Human or Bovine Thrombin Using Contrast-Enhanced Ultrasound Guidance," *The American Journal of Cardiology*, 93(12):1540-2.
 14. Vaidhyanath, R. & Blanshard, K. S. (2003). 'Treatment of a Popliteal Artery Pseudoaneurysm,' *Radiol*, 76(903):195-8.
 15. Selby, J. B. Jr, Bittner, G. M., Tegtmeyer, C. J. & Tribble, C. G. (1989). "Balloon Occlusion of a Pseudo-Aneurysm in a Below-The-Knee Amputation Stump," *American Journal of Roentgenology*. Sep; 153(3):625-7.
 16. Sugimoto, T., Kitade, T., Morimoto, N. & Terashima, K. (2004). "Pseudoaneurysms of Peroneal Artery: Treatment with Transcatheter Platinum Coil Embolization," *Annals of Thoracic and Cardiovascular Surgery*, 10(4):263-5.
 17. Edwards, H., Martin, E. & Nowygrod, R. (1982). "Nonoperative management of a Traumatic Peroneal Artery False Aneurysm," *Journal of Trauma-Injury Infection & Critical Care*; 22: 323-6.
 18. Carlin, R. E., Papenhausen, M., Farber, M. A. et al. (2001). "Sural Artery Pseudoaneurysms after Knee Arthroscopy: Treatment with Transcatheter Embolization," *Journal of Vascular Surgery*; 33:170-3.