



IBIMA
Publishing
mobile

Journal of Research in Obesity

Vol. 2014 (2014), Article ID 558641, 44 minipages.

DOI:10.5171/2014.558641

www.ibimapublishing.com

Copyright © 2014 Reetika Tripathi, QuynhDoan, Vikram Sabhaney and Kathy Boutis. Distributed under Creative Commons CC-BY 3.0

Research Article

Accuracy of the Weight-for-age Index in Identifying Obese Children in the Emergency Setting

Authors

Reetika Tripathi

University of Toronto, University Avenue, Toronto

QuynhDoan and Vikram Sabhaney

Department of Pediatrics, University of British Columbia, Division of
Emergency Medicine, Vancouver BC

Kathy Boutis

Division of Emergency Medicine, Department of Pediatrics, the Hospital for
Sick Children and the University of Toronto; University Avenue, Toronto

Received Date: 15 December 2013; Accepted Date: 6 February 2014;
Published Date: 30 April 2014

Academic Editor: Giorgio Bedogni

Cite this Article as: Reetika Tripathi, QuynhDoan, Vikram Sabhaney and Kathy Boutis (2014), "Accuracy of the Weight-for-age Index in Identifying Obese Children in the Emergency Setting," Journal of Research in Obesity, Vol. 2014 (2014), Article ID 558641, DOI: 10.5171/2014.558641

Abstract

Obesity has been defined as weight-for-age $\geq 95^{\text{th}}$ percentile for research/clinical settings like the emergency department where height is not routinely available. Our main objective was to determine the sensitivity of weight-for-age $\geq 95^{\text{th}}$ percentile in identifying obesity, using Body Mass Index (BMI)-for-age $\geq 95^{\text{th}}$ percentile as the reference standard. We also determined the specificity, and predictive values, and the correlation between weight and BMI-for-age values.

This was a cross-sectional study with prospectively collected data conducted at two urban, tertiary care pediatric emergency departments in Canada. Children between 2 and 17 years of age with acute extremity injuries were enrolled.

Of the 2259 participants, 1283 (56.1%) were male and the mean (SD) age was 9.5 (4.1) years. Using weight-for-age $\geq 95^{\text{th}}$ percentile, 326 [14.3% (12.9, 15.7)] were classified as obese, while using BMI-for-age $\geq 95^{\text{th}}$ percentile criteria determined that 363 [16.1% (14.6, 17.7)] were obese, $p < 0.0001$. The sensitivity (95% CI) of weight-for-age $\geq 95^{\text{th}}$ percentile to identify obesity was 62.2% (57.0, 67.2). The specificity, positive and negative predictive values were 94.5 % (93.6, 95.6), 14.4% (13.0, 16.0), and 85.6% (84.0, 87.0), respectively. The correlation between BMI and weight-for-age was 0.71 (0.68, 0.73).

Although there may be limited validity in using this weight-for-age cut-off as a definition for obesity in research that includes children with acute injuries, the high specificity may provide

some clinical utility in identifying two-thirds of children who are obese while minimizing false labeling of this condition.

Keywords: Obesity, Pediatrics, Definition.

Introduction

Pediatric obesity is characterized by the World Health Organization as one of the most serious public health challenges of the 21st century. (Formiguera and Canton, 2004) In 2010, an estimated 43 million preschool children world-wide were identified as overweight or obese, representing a 60% increase since 1990. (de Onis et al., 2010) Within North America rates of obesity have steadily increased, and in the Canada between 2009-2011, 31.5% of children were classified as overweight (19.9%) or obese (11.7%). (Ogden et al., 2012, Tremblay and Willms, 2000, Roberts KC et al., 2012) Children who are obese are at risk of many acute and chronic medical complications and require more medical care,(Daniels, 2006) including frequent use

of the emergency department. (Estabrooks and Shetterly, 2007, Prendergast et al., 2011)

Research and public health initiatives that examine the relationship between body habitus and a specific medical condition often uses the definition of Body Mass Index (BMI) > 95th percentile to define obesity. (Kuczmariski RJ et al., 2002) However, height is not routinely recorded in acute care clinical settings like the emergency department, and therefore weight-for-age > 95th percentile as a cut-off for defining obesity has been used in these situations. (Backstrom et al., 2012a, Leet et al., 2005a, Pomerantz et al., 2010a, Whitaker and Orzol, 2006) As such, research examining the association between weight and common urgent complaints like injuries often uses the weight-for-age > 95th percentile metric to define obesity. (Backstrom et

al., 2012b, Leet et al., 2005b, Pomerantz et al., 2010b, Ryan et al., 2010) Yet, there is currently limited evidence demonstrating how accurate the weight-for-age > 95th percentile is in identifying obesity. One study by Stettler et al used national data from the United States and determined that weight-for-age > 95th percentile was 82% sensitive in the detection of obesity relative to the BMI > 95th percentile definition. (Stettler et al., 2007) But, validating this sensitivity in children with injuries that present to the acute care setting is currently lacking.

Our main objective was to determine the accuracy of weight-for-age \geq 95th percentile in identifying the obese children who presented to the emergency department with an extremity injury relative to the BMI \geq 95th percentile reference standard. We also examined the correlation of weight-for-age and BMI-for-age, and

the distribution of differences between the weight-for-age and BMI-for-age percentiles.

Methods

Study Design and Setting

This was a cross-sectional study with prospectively collected data conducted at two urban, tertiary care pediatric emergency departments in Canada.

Study Setting and Population

Eligible children were between the ages of two and 17 years and presented to the emergency department between June 1, 2011

and December 31st, 2012 with a non-penetrating injury to an extremity. We selected extremity injuries since they one of the most common pediatric presenting complaints to emergency departments and affect all ethnic and socioeconomic backgrounds. (Pomerantz et al., 2010b) We excluded families who could not speak English, children with brittle or fragile bones, pre-existing neuromuscular disorder, global developmental delay and children with poly-trauma. The research ethics review boards of both institutions approved the study.

Definitions

Children were classified as obese in accordance with weight if they were at or above the 95th percentile for weight-for-age and gender based on data from Centers for Disease Control.

(Kuczmarski RJ et al., 2002, Ogden CL and Flegal KM, 2010) Using the BMI standard, children with a BMI-for-age at or above the 95th percentile were labelled as "obese." (Kuczmarski RJ et al., 2002, Ogden CL and Flegal KM, 2010) The weight-for-age and BMI-for-age percentiles are based on the Centre for Disease Control growth charts, (Kuczmarski RJ et al., 2002) which are based on children and adolescents who participated in national studies conducted between 1963 and 1994.

Patient Recruitment

At Site 1, patients were enrolled by on-site trained research assistants, who provided 15-hour daily research presence in the emergency department. At Site 2, there was intermittent

coverage by research assistants whereby research assistants provided coverage on average from 1000 to 2200 daily.

At both hospitals, following triage, potential study participants were identified by monitoring the site-specific emergency department tracking system for appropriate complaints/triage histories that suggested the presence of an extremity injury. Research assistants then approached families to screen for eligibility. If eligible, research assistants obtained informed consent and any applicable assent. A record of those not enrolled was kept to assess for enrollment bias. On a study form, we recorded age calculated using date of birth (day/month/year) and date of presentation (day/month/year), sex, height measured in centimeters from the top of the head to the heel of the patient, and weight measured in kilograms using a hospital

standard scale. Height was measured using a wall stadiometer in ambulating patients, and in those who were non-ambulatory height was measured using a measuring tape while lying straight on the examination bed.

Study Outcomes

The primary outcome was obesity (yes/no) as detected by BMI-for-age and the predictor is obesity (yes/no) as detected by weight-for-age. Weight-for-age and BMI-for-age percentiles were calculated using standardized methods (<http://www.cdc.gov/nccdphp/dnpa/growthcharts/training/modules/module1/text/intro.htm>) and double checked for accuracy tools available from Baylor College of Medicine and USDA/ARS Children's Nutrition Research Center, Houston

(<http://www.bcm.edu/bodycomplab/Flashapps/bmiVAgeChartpage.htm>) and the British Columbia Children's Hospital Endocrinology and Diabetes Unit, Vancouver (<http://www.bcchildrens.ca/Services/SpecializedPediatrics/EndocrinologyDiabetesUnit/ForProfessionals/AnthropometricCalculators.htm>). Secondary outcomes included a) the specificity, positive and negative predictive values of weight-for-age $\geq 95^{\text{th}}$ percentile; b) the diagnostic accuracy of the 3rd, 5th, 10th, 20th, 50th, 90th, 97th percentile weight-for-age values; c) the predictive value of patient demographic variables (age as a continuous variable and categorized in three relatively physiologically similar age groups of 2-5, 6-11, 12-17 years; sex) in accurately identifying obesity using weight-for-age $\geq 95^{\text{th}}$ percentile relative to the BMI standard; d) the correlation of weight-for-age and BMI-for-age values; and e) the proportion of patients where

weight-for-age was consistent with the BMI-for-age percentiles within a specific range of percentage points.

Sample Size Considerations

This study was powered for the primary outcome, the sensitivity of the weight-for-age > 95th percentile to identify obesity compared to BMI. Based on previous data from a similar population, we expect the prevalence of obesity to be 11% of our patient sample. (Campbell et al., 2013) Assuming the weight for age percentile method is at least 80% sensitive, (Stettler et al., 2007) in order to achieve +/- 5% precision with 95% confidence on the sensitivity, the estimated required sample size was 2235 subjects.

Statistical Analyses

Descriptive analyses were used to describe demographics. Proportions were reported with respective 95% confidence intervals. Logistic regression was used to determine if there were any significant associations between the binary outcome (accurate/not accurate) weight-for-age \geq 95th percentile and the aforementioned patient demographic variables. Odds of the outcome for a given variable were reported with respective 95% confidence intervals. Goodness-of-fit of final model to the data was tested using the Hosmer-Lemeshow test ($p > 0.05$ rejects the null hypothesis of poor fit). A receiver operator curve was plotted to check the predictive ability of the model to predict on the binary outcome of interest (accuracy of weight-for-age \geq 95th percentile to define obesity as per BMI-for-age - i.e. proportion

"accurate"). We reported the area under the receiver operator curve with respective 95% confidence interval. Pearson's correlation coefficient was used to determine the correlation of weight-for-age and BMI-for-age. Finally, we described the distribution of differences in weight-for-age and BMI-for-age percentiles, and calculated cross-classification of categories of weight-for-age and BMI-for-age percentiles. All analyses were performed using SAS statistical software (version 9.2; SAS Institute Inc, Cary, NC).

Results

Data from 2259 children were available for analysis (Figure 1). Of these, 1283 (56.1%) were male and the mean (SD) age was 9.5 (4.1) years. The mean (SD) height and weight were 38.2

(20.3) kg and 137 (27.4) cm, respectively. Using BMI-for-age criteria for obesity, 16.1% [95% CI 14.7 - 17.7] were classified as obese, while using the weight-for-age 14.3% [95% CI 13.0 - 15.9] were classified as obese, $p < 0.0001$ (Table 1).




Figure 1: Patient Enrolment

Table 1. Number of Subjects and Prevalence of Obesity Status by Sex and Age Groups

Please See Table 1 in Full PDF Version

The sensitivity of weight-for-age \geq 95th percentile to identify obese children (primary outcome) was 62.2% [95% CI 57.2-67.2] (Table 2). The specificity, positive and negative predictive values for this weight-for-age cut-off were 94.7% [95% CI 93.7-95.7], 69.3% [95% CI 64.3-74.3], and 92.9% [95% CI 91.7-94.0], respectively. There was a statistically significant trend of decreasing sensitivity for increasing weight-for-age percentile ($p < 0.001$), with the highest sensitivity of weight-for-age \leq 3% at 96.1% [95% CI 94.1-98.1] and the lowest at 51.7% [95% CI 46.6-56.9] for weight-for-age \geq 97th percentile.

Table 2. For Each Weight-For-Age Percentile Cut-Off Point, Prevalence of Subjects at or above the Cut-Off, Sensitivity, Specificity, PPV and NPV, with 95% CI's, for Obese (BMI at or above the 95th Percentile) Status in a Sample of Pediatric Patients Presenting to the Emergency Department

Please See Table 2 in Full PDF Version

Weight-for-age \geq 95th percentile did not have a higher odds of being accurate in children 6-11 versus 2-5 years [OR=0.87; 95% CI 0.48, 1.56], those 12-17 versus 2-5 years [OR=0.61; 95% CI 0.35, 1.08], or female versus male sex [OR=1.14; 95% CI 0.86, 1.49]) (Hosmer-Lemeshow $p = 0.91$ (rejecting the null hypothesis that; Predictive ability of the logistic regression model, area under the curve = 0.55; 95% CI 0.52, 0.59).

In 80% of cases, the difference between the weight-for-age and BMI-for-age percentiles was less than 5%. The BMI-for-age and weight-for-age percentiles agreed to within 0 to 2.5 percentage points for 50.3% of children. In the remaining cases, 31.9% had weight-for-age percentiles >2.5 percentage points higher than their BMI-for-age percentiles, while 17.7% had weight-for-age percentiles > 2.5 percentage points lower than their BMI-for-age percentiles (Figure 2). The Pearson's correlation between the BMI-for-age and weight-for-age was 0.71.




Figure 2. Proportion of Patients Weight-For-Age Percentiles that Were Consistent with Body Mass Index-For-Age by a Specified Range

While 69.3% of children met criteria for obesity (≥ 95 th percentile) using weight-for-age and BMI-for-age criteria, an

additional 28.4% of those who were classified as obese per weight-for-age would have still been classified as overweight (i.e. \geq 85th percentile) by BMI-for-age criteria (Table 3). Further, of the 1593 cases classified as weight-for-age \leq 84.9th percentile, 1410 (88.5%) were similarly classified by the BMI-for-age. Nevertheless, 136/362 (37.6%) who met criteria for BMI-for-age defined obesity were classified as non-obese by the weight-for-age metric, including 13% who were reported at \leq 50th weight for age percentile, accounting for the compromised sensitivity of weight-for-age.

Table 3. Cross-Classification of Categories of Weight-For-Age and Body Mass Index-For-Age Percentiles

	Body Mass Index-for-age percentile				
Weight-for-age percentile	≥ 95 N=362	90 - 94.5 N=168	85 - 89.9 N=146	50 - 84.9 N=781	≤ 50 N=797
≥ 95, no. (%)	226 (62.4)	56 (33.3)	25 (17.1)	16 (2.0)	3 (0.3)
90 - 94.9, no. (%)	40 (11.0)	45 (26.8)	42 (28.8)	62 (7.9)	5 (0.6)
85 - 89.9, no. (%)	21 (5.8)	16 (9.5)	22 (15.1)	71 (9.1)	11 (1.4)
50 - 84.9, no. (%)	28 (7.7)	46 (27.4)	51 (34.9)	489 (62.6)	257 (32.2)
≤ 50th, no. (%)	47 (13.0)	5 (3.0)	6 (4.1)	143 (18.3)	521 (65.3)

Discussion

In children who presented to an emergency department with an extremity injury, we demonstrated that the weight-for-age \geq 95th percentile had a sensitivity of 62% and a specificity of 95% in diagnosing obesity relative to the BMI reference standard.

Alternatively, the correlation between the weight-for-age versus BMI-for-age definition for obesity was very good and the majority of children had BMI-for-age and weight-for-age values that were within five percentage points.

The sensitivity of weight-for-age $>$ 95th percentile in identifying obesity relative to BMI standards is lower in this study than previously reported. A study by Stettler et al demonstrated a sensitivity of 82% in a study that included over 12,000 children

aged 2-19 years using data from the 1994-2004 the National Health and Nutrition Examination Survey in the United States. (Stettler et al., 2007) The latter report was designed to represent the entire American population, while our study was a subset of children with injuries who presented to an emergency department in Canada. One factor that may have contributed to the discrepancy in reported sensitivities is the difference in ethnicities between the studies since BMI and body fat in children differs between ethnic groups. (Deurenberg et al., 1998) In Stettler et al, one-third of the study sample was Mexican American while ethnicity data from our emergency departments demonstrate that about one-third are Southeast Asian Canadians and only 8% from Central/South America. (Boutis K et al., 2013) Further, our study population may have had a different height/weight distribution than that in the Stettler paper. That

is, we may have had a higher percentage of children that were shorter and obese with a weight-for-age percentile that did not meet the obesity cut-off. The latter may be relevant in our study since compared to the general population, there is a greater frequency of childhood obesity amongst those who present to emergency departments, (Prendergast et al., 2011) and we may have had on average shorter children given the relatively higher proportion of Southeast Asian participants. Thus, weight-for-age > 95th percentile may have falsely labelled some shorter children as non-obese, thereby compromising the sensitivity of this definition for obesity. Interestingly, despite the relatively low reported sensitivity, differences between BMI-for-age and weight-for-age percentiles were relatively small, with 80 percent of children having BMI-for-age and weight-for-age values that were within five percentage points. Further, cross-classification

of weight-for-age and BMI percentiles demonstrated that children with a weight-for-age > 95th percentile, almost 94% of these also had a BMI-for-age > 85th percentile (overweight - obese). Thus, for the majority of children the age-for-weight cut-points produce results that are similar to the BMI-for age cut-points, although these differences still resulted in many children being falsely labeled as non-obese. In this study, the specificity and negative predictive values of weight-for-age > 95th percentile in this population were very high, above 93%, and comparable to previously reported values. (Stettler et al., 2007) The latter implies that there may be some value in using weight-for-age > 95th percentile as a clinical screening tool for obesity in the emergency department. An earlier study out of the United States identified that approximately 30% of children who present to an urban emergency department are obese, (Prendergast et al.,

2011) and the reported prevalence in this study's population was approximately 16%. Thus, the emergency setting may be a significant untapped resource in identifying childhood obesity. If we use weight-for-age > 95th percentile as a screening tool, we can identify approximately two thirds of obese children without falsely labelling others as obese. Emergency departments have been shown to be a key component of the health care system by serving as connectors between acute care and preventative/outpatient management. (Prendergast et al., 2011) Thus, timely referral from the emergency department to a pediatrician and/or specialty clinics focusing on childhood obesity may be an important step in the management of this condition, especially if early intervention has any role in reducing longer-term morbidity. (O'Brien SH et al., 2004) Simultaneously, given the high specificity of the weight-for-age > 95th percentile

cut-off, we can avoid erroneous stigmatization of this label and potentially unnecessary treatments or interventions.

Specific age categories and sex were not shown to be significant predictors of the accuracy of the weight-for-age ≥ 95 percentile cut off defining obesity. Although this did not achieve significance, there was a notable trend that weight-for-age > 95 percentile in younger children (aged 2-5 years) had a lower odds (0.61) of accuracy than older (12-17 years) children. There is currently a paucity of literature available to compare our findings to other data and the rationale for this likely requires further study.

Limitations

Although BMI-for-age is considered a commonly accepted and feasible reference standard to identify pediatric obesity, it may be limited in its accuracy to diagnose obesity since this index does not directly measure body fat percentage. (Freedman and Sherry, 2009) However, recently BMI has been shown in the pediatric population to have moderately high sensitivity and positive predictive value, along with a high specificity in identifying children with excess body fatness, (Freedman and Sherry, 2009) and it is also the basis for the development of the BMI-for-age (and sex) cut points from the Centres for Disease Control, World Health Organization and the International Obesity Task Force. We did not collect data on participant ethnicity, and this would have been an important variable to collect to examine its effect

on the accuracy of weight-for-age > 95% percentile definition of obesity compared to BMI-for-age metric. Further, our study population consisted of children who presented to the emergency department with extremity injuries, and the results may not be generalizable to other populations and/or settings.

Conclusions

Although there may be limited validity in using this weight-for-age cut-off as a definition for obesity in research that includes children with acute injuries, the high specificity may provide some clinical utility in identifying two-thirds of children who are obese while minimizing false labeling of this condition.

Acknowledgements

We would like to acknowledge the Pediatric Research Academic Initiative at SickKids Emergency (PRAISE). This research would also not have been possible without the superb efforts of the program manager, Johanna Crudden and the participating PRAISE research assistants. Further, we would like to recognized the excellent efforts of the participating research assistants at British Columbia Children's Hospital Emergency Department who were responsible for the recruitment/enrolment of patients at this site.

References

Backstrom, I. C., Maclennan, P. A., Sawyer, J. R., Creek, A. T., Rue lli, L. W. & Gilbert, S. R. (2012a). "Pediatric Obesity and Traumatic Lower-Extremity Long-Bone Fracture Outcomes," *Journal of Trauma and Acute Care Surgery*, 73, 966-71.

Backstrom, I. C., Maclennan, P. A., Sawyer, J. R., Creek, A. T., Rue, L. W., 3rd & Gilbert, S. R. (2012b). "Pediatric Obesity and Traumatic Lower-Extremity Long-Bone Fracture Outcomes," *Journal of Trauma and Acute Care Surgery*, 73, 966-71.

Boutis, K., Cogollo, W., Fischer, J., Freedman, S. F., Bendavid, G. & Thomas (2013). "Parental Knowledge of Potential Cancer Risks

from Exposure to Computed Tomography," *Pediatrics*, 132, 305-11.

Campbell, J., Alqhatani, A., Mcrae, L., Kissoon, N. & Doan, Q. (2013). "Body Mass Index and the Odds of Acute Injury in Children," *Pediatric Emergency Care*, 29, 21-5.

Daniels, S. R. (2006). "The Consequences of Childhood Overweight and Obesity," *The Future of Children*, 16, 47-67.

De Onis, M., Blossner, M. & Borghi, E. (2010). "Global Prevalence and Trends of Overweight and Obesity among Preschool Children," *American Journal of Clinical Nutrition*, 92, 1257-64.

Deurenberg, P., Yap, M. & Van Staveren, W. A. (1998). "Body Mass Index and Percent Body Fat: A Meta Analysis among Different Ethnic Groups," *International Journal of Obesity & Related Metabolic Disorders*, 22, 1164-71.

Estabrooks, P. A. & Shetterly, S. (2007). "The Prevalence and Health Care Use of Overweight Children in an Integrated Health Care System," *Archives of Pediatrics & Adolescent Medicine*, 161, 222-7.

Formiguera, X. & Canton, A. (2004). "Obesity: Epidemiology and Clinical Aspects," *Best Practice & Research. Clinical Gastroenterology*, 18, 1125-46.

Freedman, D. S. & Sherry, B. (2009). "The Validity of Bmi as an Indicator of Body Fatness and Risk Among Children," *Pediatrics*, 124 Suppl 1, S23-34.

Kuczmarski, R. J., Ogden, C. I. & Guo, S. S. (2002). "2000 Cdc Growth Charts for the United States: Methods and Development," *Vital and Health Statistics*, 11, 1-190.

Leet, A. I., Pichard, C. P. & Ain, M. C. (2005a). "Surgical Treatment of Femoral Fractures in Obese Children: Does Excessive Body Weight Increase the Rate of Complications?," *The Journal of Bone and Joint Surgery*, 87, 2609-2613.

Leet, A. I., Pichard, C. P. & Ain, M. C. (2005b). "Surgical Treatment of Femoral Fractures in Obese Children: Does Excessive Body

Weight Increase the Rate of Complications?," *The Journal of Bone and Joint Surgery*, 87, 2609-13.

O'brien, S. H., Holubkov, R. & Reis, E. C. (2004). "Identification, Evaluation, and Management of Obesity in an Academic Primary Care Center," *Pediatrics*, 114, E154-9.

Ogden, C. L. & Flegal, K. M. (2010). "Changes in Terminology for Childhood Overweight and Obesity," *National Health Statistics Reports*, 25, 1-5.

Ogden, C. L., Carroll, M. D., Kit, B. K. & Flegal, K. M. (2012). "Prevalence of Obesity and Trends in Body Mass Index among Us Children and Adolescents, 1999-2010," *The Journal of the American Medical Association*, 307, 483-490.

Pomerantz, W. J., Timm, N. L. & Gittelman, M. A. (2010a). "Injury Patterns in Obese versus Nonobese Children Presenting to a Pediatric Emergency Department," *Pediatrics*, 125, 681-685.

Pomerantz, W. J., Timm, N. L. & Gittelman, M. A. (2010b). "Injury Patterns in Obese versus Nonobese Children Presenting to a Pediatric Emergency Department," *Pediatrics*, 125, 681-5.

Prendergast, H. M., Close, M., Jones, B., Furtado, N., Bunney, E. B., Mackey, M., Marquez, D. & Edison, M. (2011). "On the Frontline: Pediatric Obesity in the Emergency Department," *Journal of the National Medical Association*, 103, 922-5.

Roberts, K. C., Shields, M., De Groh, M., Aziz, A. & Gilbert, J. A. (2012). "Overweight and Obesity in Children and Adolescents:

Results from the 2009 to 2011 Canadian Health Measures Survey," *Health Reports*, 23, 37-41.

Ryan, L. M., Teach, S. J., Searcy, K., Singer, S. A., Wood, R., Wright, J. L. & Chamberlain, J. M. (2010). "Epidemiology of Pediatric Forearm Fractures in Washington," *Journal of Trauma-Injury Infection & Critical Care*, 69, S200-5.

Stettler, N., Zomorodi, A. & Posner, J. C. (2007). "Predictive Value of Weight-For-Age to Identify Overweight Children," *Obesity (Silver Spring)*, 15, 3106-12.

Tremblay, M. S. & Willms, J. D. (2000). "Secular Trends in the Body Mass Index of Canadian Children," *CMAJ*, 163, 1429-33.

Whitaker, R. C. & Orzol, S. M. (2006). "Obesity among Us Urban Preschool Children: Relationships to Race, Ethnicity, and Socioeconomic Status," *Archives of Pediatrics & Adolescent Medicine*, 160, 578-84.